



KONGERIKET NORGE
The Kingdom of Norway

Bekreftelse på patentsøknad nr
Certification of patent application no




20035228

▷ Det bekreftes herved at vedheftede dokument er nøyaktig utskrift/kopi av ovennevnte søknad, som opprinnelig inngitt 2003.11.24

▷ It is hereby certified that the annexed document is a true copy of the above-mentioned application, as originally filed on 2003.11.24

BEST AVAILABLE COPY

2005.01.19


Ellen B. Olsen
Saksbehandler



2003 -11- 24

www.patentstyret.no

Ferdig utfylt skjema sendes til adressen nedenfor. Vennligst ikke heft sammen sidene.
Vi ber om at blankettene utfylles *maskinelt* eller ved bruk av *blokkbokstaver*. Skjema for
utfylling på datamaskin kan lastes ned fra www.patentstyret.no.

Alm.tilgj. 25 MAI 2003

1a-f

KC

A blt

Søker Den som søker om patent, blir også innehaver av en eventuell rettighet. Mar fylles ut.

Foretakets navn (fornavn hvis søker er person):

Amersham Health AS

Etternavn (hvis søker er person):

☒ Kryss av hvis søker tidligere har vært kunde hos Patentstyret.

Oppgi gjerne kundennummer:

Adresse:

Nycoveien 1-2

Postboks 4220 Nydalen

Postnummer:

0401

Poststed:

Oslo

Land:

Norge

☐ Kryss av hvis flere søkere er angitt i medfølgende skjema eller på eget ark.

☐ Kryss av hvis søker(ne) utfører mindre enn 20 årsverk (se veiledning).

☐ Kryss av hvis det er vedlagt erklæring om at patentsøker(ne) innehar retten til oppfinnelsen.

Kontaktinfo Hvem skal Patentstyret henvende seg til? Oppgi telefonnummer og eventuelt referanse.

Fornavn til kontaktperson for fullmektig eller søker:

Liv-Heidi

Etternavn:

Thoresen

☒ Telefon:

2 3 1 8 5 4 7 2

Referanse (maks. 30 tegn):

PN0384-NO

☒ Evt. adresse til kontaktperson:

Amersham Health AS

Nycoveien 1-2, Postboks 4220 Nydalen

Postnummer:

0401

Poststed:

Oslo

Land:

Norge

Fullmektig Hvis du ikke har opphevet en fullmektig, kan du gå til neste punkt.

Foretakets navn (fornavn hvis fullmektig er person):

Amersham Health AS

Etternavn (hvis fullmektig er person):

☒ Kryss av hvis fullmektig tidligere har vært kunde hos Patentstyret.

Oppgi gjerne kundennummer:

Adresse:

Nycoveien 1-2

Postboks 4220 Nydalen

Postnummer:

0401

Poststed:

Oslo

Land:

Norge

Oppfinner Oppfinnere skal alltid oppgis, selv om oppfinner og søker er samme person.

Oppfinnerens fornavn:

Alan

Etternavn:

Cuthbertson

☐ Kryss av hvis oppfinner tidligere har vært kunde hos Patentstyret.

Oppgi gjerne kundennummer:

Adresse:

Amersham Health AS

Nycoveien 1-2, Postboks 4220 Nydalen

Postnummer:

0401

Poststed:

Oslo

Land:

Norge

☐ Kryss av hvis flere oppfinnere er angitt i medfølgende skjema eller på eget ark.

ADRESSE

Postboks 8160 Dep.
København 10
0033 Oslo

TELEFON

22 38 73 00
TELEFAKS
22 38 73 01

BANKGIRO

8276.01.00192
ORGANISASJONSNR.
971526167 MVA



PATENTSTYRET
Styret for det industrielle rettssvern

SØKNAD S-TAV 2

FLERE SØKERE

FLERE OPPFINNERE

PRIORITETER

VEILEDNING



7 Tittel: Gi en kort benevnelse eller tittel for oppfinnelsen (ikke over 256 tegn, inkludert mellomrom).

Tittel:
Kontrastmiddel

7 PCT: Fylls bare ut hvis denne søknaden er en videreføring av en tidligere innlevert internasjonal søknad (PCT).

PCT-søknadens dato og nummer:

PCT

/

7 Prioritetskrav: Hvis du ikke har søkt om denne oppfinnelsen tidligere (i et annet land eller i Norge), kan du gå videre til neste punkt.

Prioritet kreves på grunnlag av tidligere innlevert søknad i Norge eller utlandet:

Opplysninger om tidligere søknad. Ved flere krav skal tidligste prioritet angis her:

Inngivelsesdato (åååå.mm.dd):

Landkode:

Søknadsnummer:

☐ Flere prioritetskrav er angitt i medfølgende skjema, eller på eget ark.

7 Mikroorganisme: Fylls bare ut hvis oppfinnelsen omfatter en mikroorganisme.

Søknaden omfatter en kultur av mikroorganismer. Deponeringssted og nummer må oppgis:

☐ Prøve av kulturen skal bare utleveres til en særlig sakkyndig.

Deponeringssted og nummer (barnett gjøres eget ark).

7 Avdelt/utskilt: Hvis du ikke har søkt om patent i Norge tidligere, kan du gå videre til neste punkt.

Søknaden er avdelt eller utskilt fra tidligere levert søknad i Norge:

☐ Avdelt søknad

Informasjon om opprinnelig

Dato (åååå.mm.dd):

Søknadsnummer:

☐ Utskilt søknad

søknad/innsendt tilleggs materiale

7 Annen:

☒ Søknaden er også levert per telefaks.

Oppgi dato (åååå.mm.dd):

2 0 0 3 1 1 2 4

☐ Jeg har bedt om forundersøkelse.

Oppgi nr (årstall - nummer - bokstav):

7 Vedlegg: Angi hvilken dokumentasjon av oppfinnelsen du legger ved, samt andre vedlegg.

☒ Eventuelle tegninger i to eksemplarer

Oppgi antall tegninger: 1

☒ Beskrivelse av oppfinnelsen i to eksemplarer

☒ Patentkrav i to eksemplarer

☒ Fullmaktsdokument(er)

☐ Sammendrag på norsk i to eksemplarer

☐ Overdragelsesdokument(er)

☐ Dokumentasjon av eventuelle prioritetskrav (prioritetsbevis)

☐ Erklæring om retten til oppfinnelsen

☐ Oversettelse av internasjonal søknad i to eksemplarer (kun hvis PCT-felt over er fylt ut)

7 Dato/underskrift: Søknad du har fylt ut punktene under "Søker", "Oppfinner" og "Vedlegg", Signer søknaden.

Sted og dato (blokkbokstaver):

Oslo, 24.11.2003

Navn i blokkbokstaver:

Liv-Heidi Thoresen

Signatur:

Liv-Heidi Thoresen

NB! Søknadsavgiften vil bli fakturert for alle søknader (dvs. at søknadsavgiften ikke skal følge søknaden).
Betallingsfrist er ca. 1 måned, se faktura.



PATENTSTYRET
Styret for det industrielle rettsvern

SØKNAD s. 2 av 2

CONTRAST AGENT

1e

Field of Invention

The present invention relates to targeted contrast agents suitable for use in diagnostic imaging techniques in which a disease state may be imaged. More specifically the invention relates to contrast agents for the imaging of diseases associated with the up-regulation of the Angiotensin II type receptor AT_1 . The invention describes ligands designed to have increased potency and optimised excretion and biodistribution profiles compared to existing pharmaceutical preparations.

Diseases which can be detected by the use of a contrast agent targeting the AT_1 receptor are congestive heart failure (CHF), arteriosclerosis and fibrosis in organs like heart, lungs and liver.

Background of Invention

Angiotensin II (Ang II) - the octapeptide (Asp-Arg-Val-Tyr-Ile-His-Pro-Phe) - is a pleiotropic vasoactive peptide that binds to two distinct receptors: the Ang II type 1 (AT_1) and type 2 (AT_2) receptors. Activation of the renin-angiotensin-aldosterone system (RAAS) results in vascular hypertrophy, vasoconstriction, salt and water retention, and hypertension. These effects are mediated predominantly by AT_1 receptors. Paradoxically, other Ang II-mediated effects, including cell death, vasodilation, and natriuresis, are mediated by AT_2 receptor activation. The understanding of Ang II signalling mechanisms remains incomplete. AT_1 receptor activation triggers a variety of intracellular systems, including tyrosine kinase-induced protein phosphorylation, production of arachidonic acid metabolites, alteration of reactive oxidant species activities, and fluxes in intracellular Ca^{2+} concentrations. AT_2 receptor activation leads to stimulation of bradykinin, nitric oxide production, and prostaglandin metabolism, which are, in large part, opposite to the effects of the AT_1 receptor. (See: Berry C, Touyz R, Dominiczak AF, Webb RC, Johns DG. Am J Physiol Heart Circ Physiol. 2001 Dec;281(6):H2337-65.

Angiotensin receptors: signalling, vascular pathophysiology, and interactions with ceramide). Ang II is the active component of the renin-angiotensin-aldosterone system (RAAS). It plays an important physiological role in the regulation of blood pressure, plasma volume, sympathetic nervous activity, and thirst responses. Ang II also has a pathophysiological role in cardiac hypertrophy, myocardial infarction, hypertension, chronic obstructive pulmonary disease, liver fibrosis and atherosclerosis. It is produced systemically via the classical RAAS

and locally via tissue RAAS. In the classical RAAS, circulating renal-derived renin cleaves hepatic-derived angiotensinogen to form the decapeptide angiotensin I (Ang I), which is converted by angiotensin-converting enzyme (ACE) in the lungs to the active Ang II. Ang I can also be processed into the heptapeptide Ang-(1-7) by tissue endopeptidases.

The RAAS system is illustrated schematically in Figure 1 hereto which is based on Figure 1 in the article by Foote et al. in *Ann. Pharmacother.* **27**: 1495-1503 (1993).

In addition to the RAAS playing an important role in the normal cardiovascular homeostasis, over activity of the RAAS has been implicated in the development of various cardiovascular diseases, such as hypertension, congestive heart failure, coronary ischemia and renal insufficiency. After myocardial infarction (MI), RAAS becomes activated. Specifically the AT₁ receptor seems to play a prominent role in post-MI remodelling, since AT₁ receptor expression is increased after MI and in left ventricular dysfunction. Therefore drugs that interfere with RAAS, such as ACE inhibitors and AT₁ receptor antagonists, have been shown to be of great therapeutic benefit in the treatment of such cardiovascular disorders.

For heart, kidneys, lungs and liver alike, fibrosis represents a common pathway to their failure. Understanding pathophysiologic mechanisms involved in organ fibrosis are therefore of considerable interest, particularly given the potential for protective pharmacological strategies. Tissue repair involves inflammatory cells, including members of the monocyte/macrophage lineage, integral to initiating the repair process; and myofibroblasts, phenotypically transformed interstitial fibroblasts, responsible for collagen turnover and fibrous tissue formation. Each of these cellular events in the microenvironment of repair are associated with molecular events that lead to the de novo generation of angiotensin II (Ang II). In an autocrine/paracrine manner, this peptide regulates expression of TGF-beta 1 via angiotensin (AT₁) receptor-ligand binding. It is this cytokine that contributes to phenotypic conversion of fibroblasts to myofibroblasts (myoFb) and regulates myofibroblast turnover of collagen. Angiotensin-converting enzyme (ACE) inhibition or AT₁ receptor antagonism each prevent many of these molecular and cellular responses that eventuate in fibrosis and therefore have been found to be protective interventions.

(See: Weber KT. Fibrosis, a common pathway to organ failure: angiotensin II and tissue repair. *Semin Nephrol.* 1997 Sep;17(5):467-91 and references therein).

Ang II may regulate tissue fibrosis via the activation of mesenchymal cells. For example, Ang II stimulates the proliferation of cardiac fibroblasts *in vitro* via activation of AT₁. The presence of AT₁ receptors has also been demonstrated on cardiac fibroblasts *in vitro*. Most of the profibrotic effects of Ang II appear to be mediated via this receptor; however, increased AT₂

expression on cardiac fibroblasts has been detected in hypertrophied human heart, and the balance between the expression of these two subtypes may be critical in determining the response to Ang II.

(See: Am. J. Respir. Crit. Care Med., Volume 161, Number 6, June 2000, 1999-2004 Angiotensin II Is Mitogenic for Human Lung Fibroblasts via Activation of the Type 1 Receptor Richard P. Marshall, Robin J. McAnulty, and Geoffrey J. Laurent and references therein).

The Ang II receptors can be distinguished according to inhibition by specific antagonists. AT₁ receptors are selectively antagonized by biphenylimidazoles, such as Losartan, whereas tetrahydroimidazopyridines specifically inhibit AT₂ receptors. The AT₂ receptor may also be selectively activated by CGP-42112A. This is a hexapeptide analog of Ang II, which may also inhibit the AT₂ receptor, depending on concentration). Two other angiotensin receptors have been described: AT₃ and AT₄ subtypes.

In rodents, the AT₁ receptor has two functionally distinct subtypes, AT_{1A} and AT_{1B}, with >95% amino acid sequence homology.

The second major angiotensin receptor isoform is the AT₂ receptor. It has low amino acid sequence homology (~34%) with AT_{1A} or AT_{1B} receptors. Although the exact signaling pathways and the functional roles of AT₂ receptors are unclear, these receptors may antagonize, under physiological conditions, AT₁-mediated actions inhibiting cell growth and by inducing apoptosis and vasodilation. The exact role of AT₂ receptors in cardiovascular disease remains to be defined.

Other receptors for Ang II besides AT₁ and AT₂ are known and are generally referred to as AT_{atypical} (see Kang et al., Am. Heart J. 127: 1388-1401 (1994)).

The suppression of Ang II's effects has been used therapeutically, for example in the management of hypertension and heart failure. This has been achieved in a number of ways: by the use of renin inhibitors which block the conversion of angiotensinogen to angiotensin I (the precursor to Ang II); by the use of angiotensin converting enzyme (ACE) inhibitors that block the conversion of angiotensin I to Ang II (and also block bioconversion of bradykinin and prostaglandins); by the use of anti-Ang II-antibodies; and by the use of Ang II-receptor antagonists.

Beta blockers are most commonly used in treatment of arrhythmias. Anti-arrhythmic drugs have had limited overall success and calcium channel blockers can sometimes induce

arrhythmias. No single agent shows superiority, with the possible exception of amlodarone. Short-term anti-arrhythmic benefit has been found to be offset by, depending on the specific drug, neutral or negative effects on mortality (Sanguinetti MC and Bennett, PB: Anti-arrhythmic drug target choices and screening. *Circulation* 2003, 93(6): 491-9257-263). Clearly better anti-arrhythmic drugs are needed.

A publication in *Lancet* (Lindholm, LH et al. Effect of Losartan on sudden cardiac death in people with diabetes: data from the LIFE study. *The Lancet*, 2003, 362: 619-620) revealed that AT₁ receptor antagonists in addition of being generally favourable to patients with CHF, also reduce the incidence of sudden cardiac death. There exist a few studies showing that AT₁ antagonists have an anti-arrhythmia effect on arrhythmias induced by myocardial infarct or in reperfusion after ligation of LAD

(Harada K et al. Angiotensin II Type 1a Receptor is involved in the occurrence of reperfusion arrhythmias. *Circulation*, 1998, 97:315-317. Ozer MK et al. Effects of Captopril and Losartan on myocardial ischemia-reperfusion induced arrhythmias and necrosis in rats. *Pharmacological research*, 2002, 45 (4), 257-263

Lynch JJ et al. EXP3174, the AII antagonist human metabolite of Losartan, but not Losartan nor the Angiotensin-converting enzyme inhibitor, captopril, prevents the development of lethal ischemic arrhythmias in a canine model of recent myocardial infarction. *JACC*, 1999, 34 876-884).

It has now been found that it is possible to image Ang II receptor sites in vivo using targeted contrast agents in which the targeting binding ligand has affinity for Ang II-receptor sites. The Ang II receptors are generally accessible to such contrast agents when they are administered into the blood stream. Accordingly, using such targeted contrast agents it is possible to detect diseases and disorders such as heart failure, atherosclerosis and restricted blood flow, as well as other vascular diseases and disorders, and also to monitor the progression of treatment for such diseases and disorders.

Description of Related Art

WO 98/18496 (Nycomed Imaging AS) discloses contrast agents comprising Ang II-receptor antagonists labelled for in vivo imaging.

US patent no. 5,138,069 discloses substituted imidazoles for use as Ang II receptor blockers. Further, US patent no. 5,264,581 (Cariani) discloses radioiodinated imidazole Ang II antagonists.

Summary of the invention

When using imidazoles e.g. Losartan, as the binding ligand in a conjugate with a relatively large and bulky chelate or reporter the affinity of the ligand for the selected binding site may be negatively affected.

A problem with agents like Losartan, both the drug and the chelate conjugate, is that they excrete mainly (more than 80 %) through the liver and have an affinity (K_i) which is less avid than the natural hormone Ang II. This constitutes two problems in the use of such compositions as targeting contrast agents: Firstly a small amount of the administered composition is allowed to bind to the Ang II receptor site before it is absorbed in the liver and secondly the liver up-take results in increased background activity e.g. the background from the liver may interfere with imaging of diseased areas of the heart.

It has now been found that the introduction of an amino acid comprising biomodifier/linker between the targeting ligand and the chelate or reporter moiety can reduce the liver up-take and also increases the binding affinity to Ang II receptor sites. The biomodifier/linker may be linear or branched. Hence, compositions of matter of the present invention are useful diagnostic contrast agents for *in vivo* imaging of the mammalian body. Furthermore, the patients undergoing the imaging procedure may also be on 'sartan' drug therapy. 'Sartans' e.g. Losartan are Ang II receptor antagonists used in treatment of hypertension. A contrast agent targeting Ang II receptors will experience a competition with the treatment drug for binding to the same receptor site. It is therefore considered desirable to develop contrast agents possessing higher affinity for the AT₁ receptor than the prescribed drug to avoid deleterious competition effects of the cold drug i.e. the prescribed non-detectable 'sartan' drug.

The *in vivo* detectable moiety can be any moiety capable of detection either directly or indirectly in an *in vivo* diagnostic imaging procedure e.g. by MRI, optical imaging, scintigraphy, SPECT, PET, X-ray, ultrasound, electrical impedance or magnetometric procedures.

The compositions of matter of the present invention are useful for the *in vivo* diagnostic imaging of a range of disease states (congestive heart failure (CHF), arteriosclerosis, fibrosis in organs like heart, lungs and liver) where the up regulation of Ang II receptor sites is known to be involved.

Detailed description of the invention


In a first aspect the present invention provides a composition of matter of formula I



where V is an organic group having binding affinity for an angiotensin II receptor site, L is an amino acid comprising biomodifier or linker moiety, and R is a moiety detectable in an *in vivo* imaging procedure of a human or animal body.





The ligand V may be any organic compound having affinity for Ang II receptors. Compounds having a marked affinity for particular types of Ang II receptors, such as AT₁ or AT₂, are generally preferred. Imidazole Ang II antagonist ligands are preferred and most preferred are ligands such as Losartan, Valsartan, Candesartan and Eprosartan.

The *in vivo* detectable moiety may be a reporter chelated by a chelating agent. In one embodiment of the invention the chelating agent may be represented by an agent of formula II:




where:

each R¹, R², R³ and R⁴ is independently H or C₁₋₁₀ alkyl, C₃₋₁₀ alkylaryl, C₂₋₁₀ alkoxyalkyl, C₁₋₁₀ hydroxyalkyl, C₁₋₁₀ alkylamine, C₁₋₁₀ fluoroalkyl, or 2 or more of the R¹, R², R³ and R⁴ groups, together with the atoms to which they are attached form a carbocyclic, heterocyclic, saturated or unsaturated ring, or can represent a chelating agent given by formulas a, b, c and d.



A preferred example of a chelating agent is represented by formula e, herein denoted cPn216.



Conjugates comprising chelating agents of Formula II can be radiolabelled to give good radiochemical purity, RCP, at room temperature, under aqueous conditions at near neutral pH. An advantage of radiolabelling the conjugates at room temperature is a simplified procedure in a hospital pharmacy.

For the synthesis of chelating agents of formula II it is referred to WO 03/006070, the content of which is incorporated herein by reference.

However, the compounds defined in Formula I may also comprise chelating agents, R, as defined in WO 01/77145, Table I, pages 11-15, the content of which is incorporated herein by reference.

The reporter moieties (R) in the contrast agents of formula (I) of the invention may be any moiety capable of detection either directly or indirectly in an *in vivo* diagnostic imaging procedure.

For MR imaging the reporter will either be a non zero nuclear spin isotope (such as ^{19}F) or a material having unpaired electron spins and hence paramagnetic, superparamagnetic, ferrimagnetic or ferromagnetic properties; for optical imaging the reporter will be a light scatterer (e.g. a coloured or uncoloured particle), a light absorber or a light emitter; for magnetometric imaging the reporter will have detectable magnetic properties; for electrical impedance imaging the reporter will affect electrical impedance; and for scintigraphy, SPECT, PET, and the like, the reporter will be a radionuclide.

Stated generally, the reporter may be (1) a chelatable metal or polyatomic metal-containing ion (i.e. TcO , etc), where the metal is a high atomic number metal (e.g. atomic number greater than 37), a paramagnetic species (e.g. a transition metal or lanthanide), or a radioactive isotope, (2) a covalently bound non-metal species which is an unpaired electron site (e.g. an oxygen or carbon in a persistent free radical), a high atomic number non-metal, or a radioisotope, (3) a polyatomic cluster or crystal containing high atomic number atoms, displaying cooperative magnetic behaviour (e.g. superparamagnetism, ferrimagnetism or ferromagnetism) or containing radionuclides.

Examples of particular preferred reporter groups (R) are described in more detail below.

Chelated metal reporters are preferably chosen from the group below; ^{90}Y , $^{99\text{m}}\text{Tc}$, ^{111}In , ^{47}Sc , ^{67}Ga , ^{51}Cr , $^{177\text{m}}\text{Sn}$, ^{67}Cu , ^{167}Tm , ^{97}Ru , ^{188}Re , ^{177}Lu , ^{199}Au , ^{203}Pb and ^{141}Ce .

The metal ions are desirably chelated by chelating agents on the linker moiety. Further examples of suitable chelating agents are disclosed in US-A-4647447, WO89/00557, US-A-5367080, US-A-5364613, the content of which are incorporated herein by reference.

Methods for metallating any chelating agents present are within the level of skill in the art. Metals can be incorporated into a chelating agent by any one of three general methods:

direct incorporation, template synthesis and/or transmetallation. Direct incorporation is preferred.

Thus it is desirable that the metal ion is easily complexed to the chelating agent, for example, by merely exposing or mixing an aqueous solution of the chelating agent-containing moiety with a metal salt in an aqueous solution preferably having a pH in the range of about 4 to about 11. The salt can be any salt, but preferably the salt is a water soluble salt of the metal such as a halogen salt, and more preferably such salts are selected so as not to interfere with the binding of the metal ion with the chelating agent. The chelating agent-containing moiety is preferably in aqueous solution at a pH of between about 5 and about 9, more preferably between pH about 6 to about 8. The chelating agent-containing moiety can be mixed with buffer salts such as citrate, carbonate, acetate, phosphate and borate to produce the optimum pH. Preferably, the buffer salts are selected so as not to interfere with the subsequent binding of the metal ion to the chelating agent.

Preferred non-metal atomic reporters include radioisotopes such as ^{18}F as well as non zero nuclear spin atoms such as ^{19}F , and heavy atoms such as I.

In a further embodiment of this invention, the detectable moiety R of the composition of formula I may be a moiety useful in PET imaging. R then denotes a radioemitter with positron-emitting properties. Preferred positron emitters are the radionuclides ^{11}C , ^{18}F , ^{68}Ga , ^{13}N , ^{15}O , and ^{82}Rb . Most preferred is ^{18}F and when the radioemitter is ^{68}Ga then the preferred chelating agent is DOTA.





Non metal radionuclides such as ^{18}F may be covalently linked to the biomodifier/linker moiety by a substitution or addition reaction well known in the art. For example if the linker/biomodifier moiety L is comprised of substituents that can be chemically substituted by fluorine in a covalent bond forming reaction, such as for example, substituents containing hydroxyphenyl or p-nitrobenzoyl functionality, such substituents can be labelled by methods well known in the art with a radioisotope of fluorine.

The reporter moiety R may also represent a chromophore to be used in optical imaging procedure. By chromophore is meant a group in a composition of matter, e.g. an organic or inorganic group which absorbs and/or emits light.

By light is meant electromagnetic radiation having wavelengths from 300-1300 nm.

Chromophores having absorption and/or emission maxima in the visible to far infrared range are particularly relevant.

The invention may be exemplified by Losartan derivatives and is based on attachment of biomodifier/linker and reporter moieties to the imidazol 5-position. The principle also applies to other compounds having structural similarities, e.g. Valsartan, Candesartan and Eprosartan, possessing suitable anchoring sites in the part of the molecule corresponding to the Losartan imidazole ring.

*Losartan**Valsartan**Eprosartan**Candesartan*

One role of the biomodifier/linker moiety L is to distance the relatively bulky metal complex from the active site of the binding ligand V. The biomodifier/linker moiety can be selected to increase the binding affinity of the composition for the receptor. The biomodifier/linker moiety comprise 1-40 amino acids, and preferred 1-20 amino acids, and more preferred 1-10 amino acids and most preferred 1-5 amino acids. Further the biomodifier/linker moiety may comprise one or more dicarboxylic acid units (e.g. diglycoloyl units, glycolyl units, succinyl units), ethyleneglycol units, diamines, PEG or PEG like units or combinations of the above.

The nature of the linker group can also be used to modify the biodistribution of the resulting metal complex of the conjugate e.g. the introduction of amino acids with different properties can decrease the liver up-take.

Some of the compounds of the present invention have high affinity for the AT₁ receptor. "High affinity" refers to compounds having a $K_i \leq 5\text{nM}$ and preferably $< 0.1\text{ nM}$ and most preferred K_i 's in the pM or sub pM range calculated from competitive binding assays for AT₁ and where the K_i value was determined by competition with the known high affinity vector ¹²⁵I-Sar₁Ile₈-angiotensin II. The K_i for Ang II in this assay system is around 5 nM.

Ang II-receptor antagonists derived from the so called 'sartan' class of drugs such as Valsartan, Candesartan and Eprosartan and preferably Losartan, labelled with an imaging moiety are useful diagnostic imaging agents for in vivo imaging of a human or animal body

One preferred embodiment of the invention is the ^{99m}Tc labelled contrast agents ^{99m}Tc (Losartan-Leu-diglycolyl-cPn216), ^{99m}Tc (Losartan-Leu-Gly-diglycolyl-cPn216), ^{99m}Tc (Losartan-Leu-β-Ala-diglycolyl-cPn216) and ^{99m}Tc (Losartan-Leu-Lys(Propionyl-PEG(12)-Ac)-Diglycolyl-cPn216).

The contrast agent of formula (I) are preferably administered as a pharmaceutical formulation comprising the compound of formula (I) in a form suitable for administration to a mammal, such as a human. The administration is suitable carried out by injection or infusion of the formulation such as an aqueous solution. The formulation may contain one or more pharmaceutical acceptable additives and/or excipients e.g. buffers; solubilisers such as cyclodextrins; or surfactants such as Pluronic, Tween or phospholipids. Further, stabilisers or antioxidants such as ascorbic acid, gentisic acid or para-aminobenzoic acid and also bulking agents for lyophilisation such as sodium chloride or mannitol may be added.

The present invention also provides a pharmaceutical composition comprising an effective amount (e.g. an amount effective for enhancing image contrast in an *in vivo* imaging procedure) of a composition of general formula I or a salt thereof, together with one or more pharmaceutically acceptable adjuvants, excipients or diluents.

Viewed from a further aspect the invention provides the use of a composition of formula I for the manufacture of a contrast medium for use in a method of diagnosis involving administration of said contrast medium to a human or animal body and generation of an image of at least part of said body.

Viewed from a still further aspect the invention provides a method of generating enhanced images of a human or animal body previously administered with a contrast agent composition comprising a composition of matter as defined by formula I, which method comprises generating an image of at least part of said body.

The invention further provides a method for the monitoring of the effect of treatment of heart failure and other diseases associated with up-regulation of the AT₁ receptor.

In still another aspect the invention provides a kit for the preparation of a radiopharmaceutical composition of formula (I) comprising a ligand-chelate conjugate and a reducing agent. Preferably the reducing agent is a stannous salt. The kit may further comprise one or more stabilisers, antioxidants, bulking agents for lyophilisation and solubilisers.

The three letter abbreviations used herein for the amino acids have the following meaning:

Ala	- Alanine
Asp	- Aspartic acid
Arg	- Arginine
Glu	- Glutamic acid
Gly	- Glycine
Lys	- Lysine
Leu	- Leucine
Sar	- Sarcosine
Val	- Valine
Tyr	- Tyrosine
Ile	- Isoleucine
His	- Histidine
Pro	- Proline
Phe	- Phenylalanine
Nal	- 2- Amino-3-naphtyl propionic acid
Cha	- 2- Amino-3-cyclohexyl propionic acid

The meaning of other abbreviations used herein is as follow:

DOTA	- 1,4,7,10-tetraazacyclododecan-1,4,7,10-tetraacetic acid
PEG	- polyethyleneglycol
DIEA	- diisopropylethylamine

DPPA - diphenylphosphoryl azide

DBU - 1,8 -diazabicyclo (5,4,0) undec-7- ene

DMF - dimethyl formamide

MDP - methylene diphosphonate

TFA - trifluoroacetic acid


THF - tetrahydro furan

HATU -N-[(dimethylamino)- 1H- 1,2,3 – triazolo [4,5-b]pyridino-1-ylmethylene]-N-methylmethanaminium hexafluorophosphonate N-oxide

Fmoc - 9-fluorenylmethoxycarbonyl

General procedures

Scheme 1 shows the solid phase synthesis of Losartan linker chelating agent conjugate.



Scheme 1

K_i determination

The affinity of the compounds was determined as the dissociation constant (K_d) and measured by displacement of a radiolabelled ligand of known affinity.

The affinity of the compound for the AT_1 receptor has been determined in a competition assay using membranes from CHO cells expressing the receptor. Binding of ^{125}I -Sar $_1$ -Ile $_8$ -Angiotensin II, a ligand known to bind very efficiently to AT_1 receptors, was competed with various concentrations of the test substance. The K_i is the concentration of the competing ligand in the competition assay which would occupy 50% of the receptors if no radioligand was present. K_i is calculated using the Cheng-Prussoff equation:

$$K_i = IC_{50} / (1 + (L)/K_d)$$

Where (L) is the concentration of the radiolabelled ligand used and the K_d is the dissociation constant of the radiolabelled ligand for the receptor. IC_{50} is the concentration of competing ligand which displaces 50% of the specific binding of the radioligand. The IC_{50} value for a compound may vary between experiments depending on the radioligand concentration, whereas the K_i is an absolute value.

^{99m}Tc -Labelling protocol

A preparation is made by dissolving 0,1 mg freeze-dried cPn216 derived compound in 0,2 ml (distilled and oxygen-free) water. This solution is transferred into a 10 ml nitrogen filled vial. 0,5 ml carbonate buffer, 0,5 ml $Na^{99m}TcO_4$ solution and 0,1 ml Sn-MDP solution are added. The preparation is left at room temperature for 20 minutes.


Carbonate buffer: The carbonate buffer has a pH of 9,2 and contains 8,4 mg $NaHCO_3$ and 10,6 mg Na_2CO_3 per ml water. It is purged with nitrogen gas for at least 15 minutes before use.

$Na^{99m}TcO_4$ solution: Technetium generator (e.g. Ifetec generator) eluate, diluted to a radioactive concentration of 2 GBq/ml, oxygen free.

Sn-MDP solution: This solution contains 0,131 mg $SnCl_2 \cdot 2H_2O$ and 0,925 mg MDP (methylene diphosphonate) per ml water. The solution is made freshly before use under continuous nitrogen gas purging.


Examples

Example 1 Losartan-Leu derivatised with cPn216 via a short PEG-linker (solid phase synthesis)




All reactions were carried out in a manual nitrogen bubbler apparatus.

a) Attachment of losartan to trityl derivatised solid support




Losartan (MSD, 0.236 g, 0.558 mmol) and triethylamine (Fluka, 0.233 ml, 1.67 mmol) were added to a suspension of trityl chloride resin (Novabiochem, substitution 1.24 mmol/g, 0.300 g) in DMF (5 ml). After 4 days the resin was drained and washed. An aliquot of the resin was cleaved (dichloromethane/ TFA/ trisopropylsilane, 92.5: 5.0: 2.5, 15 min). HPLC analysis (column Phenomenex Luna C18(2) 3 μ m 4.6 x 50 mm, solvents: A = water/0.1% TFA and B = acetonitrile/0.1% TFA; gradient 10-40% B over 10 min; flow 2.0 ml/min, UV detection at 214 and 254 nm) gave a peak with t_R 6.7 minutes corresponding to losartan. The resin was treated with dichloromethane/ methanol/ diisopropylethylamine solution (17 : 2 : 1, 20 ml, 1 h), washed with dichloromethane and dried.

b) Replacement of the hydroxyl group by azide



c) Reduction of the azide group to amine



Fmoc-Leu-OH (Novabiochem, 0.030 g, 0.084 mmol) and Fmoc-amino PEG diglycolic acid (Polypure, 0.045 mg, 0.084 mmol) were successively coupled to an aliquot of the resin bound amino-losartan from c) (0.042 mmol) in DMF using standard coupling reagents (HATU and DIEA) and standard Fmoc-cleavage protocol (20% piperidine in DMF). Completion of couplings were checked by standard Kaiser test. The second diglycoloyl unit was introduced using diglycolic anhydride (Aldrich, 0.010 g, 0.084 mmol) and DIEA (0.014 ml, 0.084 mmol). To the resin (containing a terminal carboxyl function) was added the chelate cPn216 (0.029 g, 0.084 mmol), PyAOP (Applied Biosystems, 0.022 g, 0.042 mmol) and DIEA (0.014 ml, 0.084 mmol). After two hours the compound was cleaved off the resin (dichloromethane/

The compound was tested *in vitro* for binding to angiotensin-II receptor AT1 (Ki 0.5 nM).

Examples 2-18 Amino acid substituted losartan derivatives

(X1)

Table 1

AA1	AA2	MW (exact)	MH ⁺ observed	Ki (nM)
Lys		549.27	550.2	55
Glu		550.22	551.1	14
Leu		534.26	535.1	0.41
D-Lys		549.27		64
D-Glu		550.22		73
D-Leu		534.26		28

Leu	Lys	662.36		0.38
Leu	Glu	663.30		
Leu	Leu	647.35		0.63
D-Leu	D-Lys	662.36		1.6
D-Leu	D-Glu	663.30		4.5
D-Leu	D-Leu	647.35		0.53
Leu	D-Leu	647.35		4
D-Leu	Leu	647.35		0.5
Phe		568.25		
Nal		618.26		46
Cha		574.29		6.9

Examples 19-25: Losartan-Leu derivatives

Compounds of the general formula (X2) were synthesised on solid support as described in Example 1 (Table 2)












Table 2

X	Abbr	MW (exact)	MH ⁺ obs.	Ki (nM)
	Losartan-Leu- Biotin	760.34	761.6	0.16
	Losartan-Leu- Diglycoloyl- cPn216	975.56	975.9	0.0006
	Losartan-Leu- Diglycoloyl- PEG(4)- Fluorescein	1182.46	1183.4	4.7
	Losartan-Leu- Propionyl- PEG(12)- Fluorescein	1491.66	1492.6	7.9
	Losartan-Leu- Gly-Diglycoloyl- cPn216	1032.58	1033.7	0.003
	Losartan-Leu-β- Ala-Diglycoloyl- cPn216	1046.59	1047.5	0.03
	Losartan-Leu- Lys(Propionyl- PEG(12)-Ac)- Diglycoloyl- cPn216	1745.01	1745.6	0.49



Claims


1. A contrast agent of formula I



where V is an organic group having binding affinity for an angiotensin II receptor site, L is a linear or branched amino acid comprising biomodifier or linker moiety, and R is a reporter moiety detectable in *in vivo* imaging of a human or animal body.

2. A contrast agent according to claim 1 where V is an imidazole-containing Angiotensin II antagonist.
3. A contrast agent according to claim 1 or 2 where V is Losartan, Valsartan, Candesartan or Eprosartan.
4. A contrast agent according to claims 1-3 where L is comprising 1-40 amino-acid residues.
5. A contrast agent according to claims 1-4 where L is comprising 1-20 amino-acid residues.
6. A contrast agent according to claims 1-5 where L is comprising 1-10 amino-acid residues.
7. A contrast agent according to claims 1-6 where L is comprising 1-5 amino-acid residues
8. A contrast agent according to claim 1-7 where L additionally comprises one or more dicarboxylic acid units, ethyleneglycol units or PEG-like components or combinations of the above.
9. A contrast agent according to claims 1-7 where L additionally comprises one or more diclycolyl, glycolyl or succinyl units or combinations of the above.
10. A contrast agent according to any of the preceding claims where L is branched.

11. A contrast agent according to claim 1-10 where R is a reporter chelated by a chelating agent.
12. A contrast agent according to any of the preceding claims wherein the chelated reporter is radionuclides, paramagnetic metal ions, fluorescent metal ions, heavy metal ions or cluster ions.
13. A contrast agent according to any of the preceding claims wherein the chelated reporter moiety comprises ^{90}Y , $^{99\text{m}}\text{Tc}$, ^{111}In , ^{47}Sc , ^{67}Ga , ^{51}Cr , $^{177\text{m}}\text{Sn}$, ^{67}Cu , ^{167}Tm , ^{97}Ru , ^{188}Re , ^{177}Lu , ^{199}Au , ^{203}Pb , ^{141}Ce or ^{18}F .
14. A contrast agent according to claim 1-13 where the chelating agent is of formula II




where:

each R^1 , R^2 , R^3 and R^4 is independently an R group;

each R group is independently H or C_{1-10} alkyl, C_{3-10} alkylaryl, C_{2-10} alkoxyalkyl, C_{1-10} hydroxyalkyl, C_{1-10} alkylamine, C_{1-10} fluoroalkyl, or 2 or more R groups, together with the atoms to which they are attached form a carbocyclic, heterocyclic, saturated or unsaturated ring.

15. A contrast agent according to any of the preceding claims where the chelating agent is of formula (e).



16. A contrast agent according to claims 1-13 where the chelating agent is DOTA.
17. A contrast agent as claimed in claims 1-16 wherein the reporter moiety is ^{99m}Tc or an iodine isotope for SPECT imaging or an ^{11}C , ^{18}F or ^{68}Ga isotope for PET imaging procedures.
18. A contrast agent according to any of the preceding claims characterised in that it is ^{99m}Tc (Losartan-Leu-diglycolyl-cPn216).
19. A contrast agent according to any of the preceding claims 1-17 characterised in that it is ^{99m}Tc (Losartan-Leu-Gly-diglycolyl-cPn216).
20. A contrast agent according to any of the preceding claims 1-17 characterised in that it is ^{99m}Tc (Losartan-Leu- β -Ala-diglycolyl-cPn216).
21. A contrast agent according to any of the preceding claims 1-17 characterised in that it is ^{99m}Tc (Losartan-Leu-Lys(Propionyl-PEG(12)-Ac)-Diglycoloyl-cPn216).
22. A pharmaceutical composition comprising an effective amount of a compound of general formula I or a salt thereof, together with one or more pharmaceutically acceptable adjuvants, excipients or diluents for use in enhancing image contrast in *in vivo* imaging.
23. Use of a compound of formula I for the manufacture of a contrast agent for use in a method of diagnosis.
24. A method of generating enhanced images of a human or animal body previously administered with a contrast agent composition comprising a compound as defined by formula I, which method comprises generating an image of at least part of said body.
25. A method of monitoring the effect of treatment of heart failure and other diseases associated with up-regulation of the AT_1 receptor.
26. A kit for the preparation of a radiopharmaceutical composition of formula I comprising a ligand-chelate conjugate and a reducing agent.

27. Kit of claim 26 where the reducing agent is a stannous salt.

28. Kit of claim 26 and 27 additionally comprising one or more stabilisers, antioxidants, bulking agents for lyophilisation and solubilisers.






Abstract

A contrast agent of formula I

V - L - R (I)

where V is an organic group having binding affinity for an angiotensin II receptor site, L is a linear or branched amino acid comprising biomodifier or linker moiety, and R is a reporter moiety detectable in *in vivo* imaging of a human or animal body.





Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/NO04/000358

International filing date: 23 November 2004 (23.11.2004)

Document type: Certified copy of priority document

Document details: Country/Office: NO
Number: 20035228
Filing date: 24 November 2003 (24.11.2003)

Date of receipt at the International Bureau: 25 January 2005 (25.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.